


Assessment



Student Information

Date of Measurement	Age (mos)	Height (in)	Weight (lbs)	Weight (ozs)	BMI	BMI Percentile
10-15-1998	66.82	47	55		17.51	90.06
11-15-1999	79.83	49.5	58		16.64	76.27
12-15-2000	92.84	50.5	65		17.92	83.58
11-15-2001	103.84	52	66		17.16	68.19
01-15-2003	117.84	55.25	87.5		20.15	87.02
09-26-2003	126.18	57.75	129		27.19	98.21

Impact of medication

The graph depicts the BMI-for-age of a female 5 years 5 months old to 10 years 5 months old. This girl was diagnosed with ADHD (attention deficit hyperactivity disorder) in 1999 and was periodically on Ritalin for 2 years. In 2002, she was diagnosed with bi-polar disorder and started on Risperidone. During the next 14 months she gained 21.5 pounds with an increase in height of 3.25 inches. Her BMI increased by 2.99 units. The next year her medication was changed to Quetiapine fumarate. Within eight months she experienced an additional weight gain of 41.5 pounds with an increase in BMI of 7.04 units. At the 98th percentile, she was now in the overweight category. *A possible change in medication and/or lifestyle counseling in nutrition and physical activity may have helped to ameliorate the known impact of these medications on weight.*